

DOCTOR RECOMMENDED  
#1

Children's  
**TYLENOL**



8 HR FAST FEVER + PAIN RELIEF

Children's  
**Motrin**



SimpleMeasure™

IMPROVED  
Dosing Accuracy

## FEVER AND DOSAGE DIARY

Use this chart to keep track of your child's fever and medicine dosing.  
Make sure any other caregivers use it as well.

Date						
Time						
Temperature						
Medicine						
Dose						
Given By						

IMPORTANT: Give your child only ONE medicine at a time containing acetaminophen.

JOIN TODAY  
**HEALTHY ESSENTIALS™**.ca

Johnson & Johnson PHARMACEUTICALS COMPANY

For more helpful advice  
visit us on



†Reduces fever for up to 8 hours.  
© Johnson & Johnson Inc. 2016



## DOSAGE GUIDE FOR FEVER & PAIN

RECOMMENDATION FOR:

Age	Weight (Note: For Most Accurate Dosing, Dose By Weight)							
	lbs	kgs	<b>Infants'</b> SUSPENSION CONCENTRATED DROPS 80 mg/1 mL 	<b>Children's</b> LIQUID SUSPENSION 160 mg/5 mL 	<b>Junior</b> STRENGTH CHILDREN'S CHEWABLES 160 mg per tablet 	<b>Infants'</b> SUSPENSION CONCENTRATED DROPS 40 mg/1 mL 	<b>Children's</b> LIQUID SUSPENSION 100 mg/5 mL 	
<b>SINGLE ORAL DOSE</b>								
<input type="checkbox"/> 0-3 mos	Under 12	Under 5.5	Consult a doctor for dosing under 4 months					
<input type="checkbox"/> 4-11 mos	12-17	5.5-7.9	1 mL	Use Infants' TYLENOL® Suspension Concentrated Drops			1.25 mL	Use Infants' MOTRIN® Suspension Concentrated Drops
<input type="checkbox"/> 12-23 mos	18-23	8-10.9	1.5 mL				1.875 mL	
<input type="checkbox"/> 2-3 yrs	24-35	11-15.9	—	5 mL	1 tablet	—	5 mL	
<input type="checkbox"/> 4-5 yrs	36-47	16-21.9	—	7.5 mL	1 ½ tablets	—	7.5 mL	
<input type="checkbox"/> 6-8 yrs	48-59	22-26.9	—	10 mL	2 tablets	—	10 mL	
<input type="checkbox"/> 9-10 yrs	60-71	27-31.9	—	12.5 mL	2 ½ tablets	—	12.5 mL	
<input type="checkbox"/> 11 yrs	72-95	32-43.9	—	15 mL	3 tablets	—	15 mL	

Infants', Children's & Junior Strength Children's TYLENOL® products dosage guide is based on a 10-15 mg acetaminophen/kg/dose. Single dose may be repeated every 4-6 hours, as needed. It is hazardous to exceed 5 doses of acetaminophen per day. Refer to package insert for dosing instructions. Infants' and Children's MOTRIN® product single doses may be given every 6 to 8 hours as needed, but do not exceed 4 doses per day unless advised by your doctor. Refer to package insert for dosing instructions.

Only Infants' MOTRIN® Suspension Concentrated Drops 40 mg/mL and Infants' TYLENOL® Suspension Concentrated Drops 80 mg/mL are labelled for children under 2 years of age. Consumer labelling for Children's MOTRIN® Suspension 100 mg/5 mL, Children's TYLENOL® Liquid Suspension 160 mg/5 mL or for Junior Strength TYLENOL FASTMELTS® Tablets 160 mg does not offer dosing for children under 2 years of age. Package label does not provide dosing for use in infants less than 4 months of age for Infants' TYLENOL® Suspension Concentrated Drops 80 mg/mL or Infants' MOTRIN® Suspension Concentrated Drops 40 mg/mL. Please consult a doctor for dosing under 4 months of age.

Infants' and Children's TYLENOL® products are indicated for the temporary relief of mild to moderate pain in a wide variety of conditions involving musculoskeletal pain, headache pain, earache, the symptomatic reduction of fever due to the common cold, flu and other viral or bacterial infections, immunization and teething pain.

Infants' and Children's MOTRIN® products are indicated for the temporary relief of minor aches and pains in muscles, bones and joints, headache, fever, the aches and fever due to the common cold or flu, immunizations, toothache (dental pain), sore throat and earache.

For more information please consult [www.tylenol.ca](http://www.tylenol.ca) or the package inserts.